



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

XXI International Grassland Congress / VIII
International Rangeland Congress

Effects of Animal Grazing on Quality of Grass and Forb Forages in Mazandaran Province of Iran

Ghasem Ali Dianati Tilaki
Tarbiat Modares University, Iran

Follow this and additional works at: <https://uknowledge.uky.edu/igc>

 Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/10-2/23>

The XXI International Grassland Congress / VIII International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Effects of animal grazing on quality of grass and forb forages in Mazandaran province of Iran

Ghasem Ali Dianati Tilaki

Assistant professor of Tarbiat Modares University, Faculty of Natural Resources and Marine Sciences, Department of Range Management, P.O.Box 46414-356, Noor-Iran. (dianatitilaki@yahoo.com)

Abstract Effects of light (3 ha/cow/yr) and moderate (1 ha/cow/yr) stocking rates on forage quality of ... were evaluated. Crude protein of forbs were generally greater than for others species under both, moderate and light stocking. Moderate stocking during rapid plant growth could positive affect on quality of available forage plants.

Key words : crude protein, dry mater digestibility, stocking rate, forage quality

Introduction Rangeland livestock productivity is influenced by the quality of forage consumed which, in turn, is a function of the quantity and quality of forage produced. Although climate is a major factor affecting forage quality in grasslands (Lauenroth et al. 1994), the major factors within a given climatic regime are the inherent productivity potential of the site and the kinds and number of plants present (i.e., range condition) (Bartolome et al. 1992). The objective of this study was to evaluate effects of moderate and light stocking rates on the nutritional quality of available forages in a northern rangeland of Iran.

Materials and methods The study was conducted at the Chahardangeh Sari in a northern mixed-grass prairie of Iran. Vegetation is composed of Grazing treatments were year-long, continuously stocked with cattle at light (3 ha/cow/yr; LG) and moderate rates (1 ha/cow/yr; MG). These treatments were initiated in 2004 and replicated twice in 4 pastures of similar conditions (Heitschmidt et al. 1991). Crude protein (CP), dry mater digestibility (DMD) and acide detergent fiber (ADF) responses of four species of grasses and three species of forbs to stocking rates were evaluated. Mean separations were accomplished using Tukey's procedure ($p < 0.05$).

Results Moderate stocking produced increases in CP and DMD for both, grasses and forbs, relative to light stocking. Forbs generally had greater forage quality than grasses in both stocking treatments (Table 1A-1B).

Table 1 Comparison between forage quality of grasses (A) and forbs (B) in different stocking rates.

(A)			(B)		
Category	Treatments		Category	Treatments	
	Light grazing (LG)	Moderate grazing (MG)		Light grazing (LG)	Moderate grazing (MG)
CP (%)	7.2 ± 0.4 ^b	8.5 ± 0.3 ^a	CP (%)	10.2 ± 1.3 ^b	11.8 ± 0.9 ^a
DMD (%)	49 ± 0.6 ^b	53.6 ± 0.4 ^a	DMD (%)	62.3 ± 3.4 ^b	68.6 ± 2.2 ^a
ADF (%)	46.3 ± 0.9 ^b	44.5 ± 0.8 ^a	ADF (%)	31.3 ± 1.4 ^b	29.2 ± 0.9 ^a

a, b significant difference ($p < 0.05$).

Discussion and conclusion Moderate stocking rate, during rapid plant growth, can positive affect forage quality of dominant grass and forb species on mixed-grass prairie of northern Iran. In addition, grass species in this study generally had lower forage quality than forb species which, is similar to findings from other parts of the world (Norton 1982, Smith et al. 1986, Holechek et al. 2005).

References

- Bartolome, J.W., and M.P. McClaran, 1992, Composition and production of California oak savanna seasonally grazed by sheep. *J. Range Mgt* 45(1):103-107.
- Heitschmidt, R.K., and J.W. Stuth (Eds). (1991) *Grazing management: An ecological perspective*. Timber press, Portland, OR 259pp.
- Laurenroth, W.K., D.G. Milchunas, J.L. Dodd, R.H. Hart, R.K. Heitschmidt, and L.R. Rittenhouse, 1994, Effects of grazing on ecosystems on the great plains. In M. Varra, W. Laycock, and R.D. Pieper (Eds), *Ecological implications of livestock herbivory in the west*. Soc. Range Mgt., Denver, Co. PP. 69-100.
- Smith, B., Leung, P.S., Love, G., 1986, Intensive grazing management: forages, animals, Men, profits. *The graziers, Honolulu, Hawaii*, 350pp.
- Norton, B.W., 1982, Differences between species in forage quality, *Farnham Royal, UK*, pp 80-110.
- Holechek, J.L., Pieper, R.D., Herbel, C.H., 2005, Range management principles and practices. *Prentice Hall, Englewood cliff*, pp 174-283.